

Fill out ALL lined sections of the form.



For use with adrenaline (epinephrine) autoinjectors

Include a colour photo if you have one OR if the student is currently enrolled, the school will add the photo.

Name: _____
Date of birth: _____

Allergen(s) MUST be identified

Confirmed allergens: _____

Family/emergency contact name(s): _____

Work Ph: _____
Home Ph: _____
Mobile Ph: _____

Plan prepared by medical or nurse practitioner: _____

I hereby authorise medications specified on this plan to be administered according to the plan
Signed: _____

Date: _____

Action Plan due for review - date: _____

Doctor to complete and sign this section

How to give EpiPen® adrenaline (epinephrine) autoinjectors

- Form fist around EpiPen® and PULL OFF BLUE SAFETY RELEASE
- Hold leg still and PLACE ORANGE END against outer mid-thigh (with or without clothing)
- PUSH DOWN HARD until a click is heard or felt and hold in place for 3 seconds REMOVE EpiPen®

EpiPen® is prescribed for children over 20kg and adults. EpiPen®Jr is prescribed for children 10-20kg

© ASCIA 2018 This plan was developed as

SIGNS OF MILD TO MODERATE ALLERGIC REACTION

- Swelling of lips, face, eyes
- Tingling mouth
- Hives or welts
- Abdominal pain, vomiting (these are signs of anaphylaxis for insect allergy)

ACTION FOR MILD TO MODERATE ALLERGIC REACTION

- For insect allergy - flick out sting if visible
- For tick allergy seek medical help or freeze tick and let it drop off
- Stay with person and call for help
- Locate adrenaline autoinjector
- Give other medications (if prescribed) _____
- Phone family/emergency contact

Mild to moderate allergic reactions (such as hives or swelling) may not always occur before anaphylaxis

WATCH FOR ANY ONE OF THE FOLLOWING SIGNS OF ANAPHYLAXIS (SEVERE ALLERGIC REACTION)

- Difficult/noisy breathing
- Difficulty talking and/or hoarse voice
- Swelling/tightness in throat
- Persistent dizziness or collapse
- Wheeze or persistent cough
- Pale and floppy (young children)

ACTION FOR ANAPHYLAXIS

- 1 Lay person flat - do NOT allow them to stand or walk
- If unconscious, place in recovery position
- If breathing is difficult allow them to sit
 - 2 Give adrenaline autoinjector
 - 3 Phone ambulance - 000 (AU) or 111 (NZ)
 - 4 Phone family/emergency contact
 - 5 Further adrenaline doses may be given if no response after 5 minutes
 - 6 Transfer person to hospital for at least 4 hours of observation
- If in doubt give adrenaline autoinjector
Commence CPR at any time if person is unresponsive and not breathing normally

ALWAYS give adrenaline autoinjector FIRST, and then asthma reliever puffer if someone with known asthma and allergy to food, insects or medication has SUDDEN BREATHING DIFFICULTY (including wheeze, persistent cough or hoarse voice) even if there are no skin symptoms
Asthma reliever medication prescribed: Y N

- If adrenaline is accidentally injected (e.g. into a thumb) phone your local poisons information centre.
- Continue to follow _____ person with the allergic reaction

For tick allergy, tick the appropriate box.

Doctor to tick relevant box.

If applicable, include prescribed medication